

Physicians Laboratory Services Cumulative Antimicrobial Susceptibility Report
All Clients, Urine Isolates | January 2024 - December 2024

| % Susceptible Gram Positive | Total* | Penicillin(2) | Ampicillin | Oxacillin(1) | Amoxicillin/Clavulanate | Cefotaxime(2) | Ceftriaxone(2) | Meropenem | Gentamicin | Ciprofloxacin(7) | Levofloxacin(7) | Moxifloxacin | Trimethoprim/Sulfa | Clindamycin(5) | Daptomycin(6) | Erythromycin(5) | Nitrofurantoin(3) | Linezolid(4) | Vancomycin | Tetracycline(7) | Gentamicin Synergy(8) | Streptomycin Synergy(8) | Penicillin (non-CSF)(2) | Cefotaxime (non-CSF)(2) | Ceftriaxone (non-CSF)(2) | |
|---|--------|---------------|------------|--------------|-------------------------|---------------|----------------|-----------|------------|------------------|-----------------|--------------|--------------------|----------------|---------------|-----------------|-------------------|--------------|------------|-----------------|-----------------------|-------------------------|-------------------------|-------------------------|--------------------------|-----|
| <i>Staph. aureus</i> (MSSA) 66% | 119 | -- | -- | 100 | -- | -- | -- | -- | 98 | 83 | 84 | -- | 100 | -- | 100 | -- | 100 | 100 | 100 | 98 | -- | -- | -- | -- | -- | |
| <i>Staph. aureus</i> (MRSA) 34% | 60 | -- | -- | 0 | -- | -- | -- | -- | 100 | 12 | 13 | -- | 90 | -- | 100 | -- | 98 | 100 | 100 | 88 | -- | -- | -- | -- | -- | |
| <i>Staph.</i> spp. (coagulase-negative) | 656 | -- | -- | 48 | -- | -- | -- | -- | 95 | 74 | 75 | -- | 74 | -- | 100 | -- | 100 | 100 | 100 | 86 | -- | -- | -- | -- | -- | |
| <i>Strep. pneumoniae</i> | 5 | 80 | -- | -- | 100 | 100 | 100 | -- | -- | -- | 100 | -- | 80 | -- | -- | -- | -- | -- | 100 | 100 | -- | -- | 100 | 80 | 100 | 100 |
| <i>Enterococcus faecalis</i> | 1177 | 100 | 100 | -- | -- | R | R | -- | -- | 81 | 90 | -- | R | R | 100 | -- | 100 | 99 | 100 | 27 | -- | -- | -- | -- | -- | |
| <i>Enterococcus faecium</i> | 95 | 24 | 26 | -- | -- | R | R | -- | -- | 21 | 28 | -- | R | R | 78 | -- | 76 | 100 | 72 | 27 | -- | -- | -- | -- | -- | |
| <i>Enterococcus</i> spp. | 13 | 92 | 100 | -- | -- | R | R | -- | -- | 77 | 100 | -- | R | R | 100 | -- | 92 | 100 | 75 | 31 | -- | -- | -- | -- | -- | |

The % susceptible for each organism/antimicrobial combination was generated by including the first isolate of that organism encountered on a given patient.

R Organism has intrinsic resistance to this antimicrobial.

- Not tested/indicated for organism.

* Antibiograms created for organisms with less than 30 total isolates have questionable statistical significance. Interpret data with caution.

(1) Oxacillin predicts susceptibility to most cephalosporins, carbapenems, and beta-lactam/beta-lactamase inhibitors.

(2) For *S. pneumoniae*: cefotaxime, ceftriaxone, and penicillin % susceptible listed is based on parenteral CSF (meningitis), parenteral non-CSF, and oral MIC breakpoints.

(3) Nitrofurantoin is reported for isolates from urine only.

(4) Linezolid is reported for isolates from non-urine sources only.

(5) For *Staphylococcus* spp.: clindamycin and erythromycin are reported for isolates from non-urine sources only.

(6) For *Staphylococcus* spp.: daptomycin is reported for isolates from non-respiratory sources only.

(7) For all *Enterococcus* spp.: ciprofloxacin, levofloxacin, and tetracycline are reported for isolates from urine only.

(8) For *E. faecalis* and *E. faecium*: gentamicin and streptomycin high-level resistance testing is reported for isolates from blood cultures only.

Physicians Laboratory Services Cumulative Antimicrobial Susceptibility Report
All Clients, Urine Isolates | January 2024 - December 2024

| % Susceptible Gram Negative | Total* | Ampicillin | Ampicillin/Sulbactam | Piperacillin/Tazobactam | Cefazolin | Ceftazidime | Ceftazidime/Avibactam | Ceftriaxone | Cefepime | Cefoxitin | Aztreonam | Ertapenem | Imipenem | Meropenem | Amikacin | Gentamicin | Tobramycin | Ciprofloxacin(3) | Levofloxacin(3) | Trimethoprim/Sulffa | Nitrofurantoin(1) | Tetracycline(2) | Tigecycline | |
|--|--------|-----------------------------|----------------------|-------------------------|-----------|-------------|-----------------------|-------------|----------|-----------|-----------|-----------|----------|-----------|----------|------------|------------|------------------|-----------------|---------------------|-------------------|-----------------|-------------|----|
| <i>Escherichia coli</i> | 7920 | 61 | 66 | 98 | 74 | 91 | 100 | 92 | 92 | 96 | 95 | 100 | 100 | 100 | 100 | 93 | 93 | 86 | 87 | 79 | 98 | 78 | 100 | |
| <i>Klebsiella pneumoniae</i> | 1620 | R | 85 | 99 | 88 | 94 | 100 | 94 | 94 | 95 | 95 | 100 | 100 | 100 | 100 | 97 | 97 | 96 | 98 | 89 | 50 | 84 | 99 | |
| <i>Proteus mirabilis</i> | 809 | 85 | 91 | 100 | 75 | 98 | 100 | 98 | 98 | 99 | 99 | 100 | -- | 100 | 100 | 92 | 92 | 82 | 89 | 83 | R | R | -- | |
| <i>Klebsiella (Enterobacter) aerogenes</i> | 252 | R | R | 88 | R | 78 | 100 | 73 | 97 | R | 86 | 96 | 68 | 100 | 100 | 98 | 99 | 99 | 99 | 97 | 28 | 91 | 100 | |
| <i>Enterobacter cloacae</i> complex | 329 | R | R | 78 | R | 72 | 100 | 62 | 87 | R | 70 | 82 | 95 | 100 | 100 | 97 | 97 | 96 | 98 | 88 | 32 | 88 | 98 | |
| <i>Serratia marcescens</i> | 85 | R | R | 82 | R | 75 | 100 | 74 | 96 | R | 76 | 98 | -- | 100 | 100 | 100 | 99 | 94 | 99 | 95 | R | 12 | 96 | |
| <i>Pseudomonas aeruginosa</i> | 564 | R | R | 96 | R | 98 | 99 | R | 97 | R | 92 | R | 82 | 93 | 98 | 88 | 99 | 90 | 91 | R | -- | R | -- | |
| <i>Acinetobacter baumannii</i> | 2 | R | 100 | -- | R | 100 | -- | 100 | 100 | R | R | R | 100 | 100 | 100 | 100 | 100 | 100 | 100 | -- | 100 | -- | -- | |
| <i>Citrobacter freundii</i> complex | 286 | R | R | 93 | R | 78 | 100 | 72 | 94 | R | 78 | 99 | 98 | 100 | 100 | 94 | 96 | 94 | 95 | 88 | 92 | 84 | 99 | |
| <i>Morganella morganii</i> | 72 | R | 6 | 97 | R | 83 | 100 | 87 | 97 | 85 | 94 | 100 | 8 | 100 | 100 | 87 | 90 | 79 | 85 | 67 | R | R | 0 | |
| <i>Providencia</i> spp. | 60 | R | 43 | 98 | R | 68 | 97 | 90 | 98 | 93 | 88 | 98 | 60 | 98 | 100 | 72 | 72 | 70 | 75 | 80 | R | R | 0 | |
| <i>Salmonella</i> spp. | 0 | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | |
| <i>Shigella</i> spp. | 0 | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | |
| <i>Stenotrophomonas</i> | 11 | R | R | R | R | 64 | -- | R | -- | R | R | R | R | R | R | R | R | R | -- | 91 | 100 | -- | R | -- |
| <i>Klebsiella oxytoca</i> | 319 | R | 64 | 96 | 14 | 92 | 100 | 90 | 92 | 97 | 90 | 100 | 100 | 100 | 100 | 94 | 94 | 96 | 98 | 87 | 93 | 87 | 100 | |
| <i>Haemophilus influenzae</i> | 1 | Beta-lactamase positive: 0% | | | | | | | | | | | | | | | | | | | | | | |

The % susceptible for each organism/antimicrobial combination was generated by including the first isolate of that organism encountered on a given patient.

R Organism has intrinsic resistance to this antimicrobial.

- Not tested/indicated for organism.

* Antibiograms created for organisms with less than 30 total isolates have questionable statistical significance. Interpret data with caution.

(1) Nitrofurantoin is reported for isolates from urine only.

(2) For *A. baumannii*: tetracycline is reported for isolates from urine only.

(3) Ciprofloxacin/Levofloxacin: Enterobacteriales & *P. aeruginosa* %Susceptible utilizes CLSI M100 28th Edition breakpoints.