

Physicians Laboratory Services Cumulative Antimicrobial Susceptibility Report
Hospital/Surgical Clients, All Isolates | January 2024 - December 2024

% Susceptible Gram Positive	Total*	Penicillin(2)	Ampicillin	Oxacillin(1)	Amoxicillin/Clavulanate	Cefotaxime(2)	Ceftriaxone(2)	Meropenem	Gentamicin	Ciprofloxacin(7)	Levofloxacin(7)	Moxifloxacin	Trimethoprim/Sulfa	Clindamycin(5)	Daptomycin(6)	Erythromycin(5)	Nitrofurantoin(3)	Linezolid(4)	Vancormycin	Tetracycline(7)	Gentamicin Synergy(8)	Streptomycin Synergy(8)	Penicillin (non-CSF)(2)	Penicillin V (oral)(2)	Cefotaxime (non-CSF)(2)	Ceftriaxone (non-CSF)(2)
<i>Staph. aureus</i> (MSSA) 75%	153	--	--	100	--	--	--	--	99	92	94	98	100	72	100	65	100	100	100	93	--	--	--	--	--	--
<i>Staph. aureus</i> (MRSA) 25%	50	--	--	0	--	--	--	--	96	32	32	76	98	74	100	20	100	100	100	88	--	--	--	--	--	--
<i>Staph. spp.</i> (coagulase-negative)	105	--	--	50	--	--	--	--	95	72	72	91	71	61	100	52	100	100	100	84	--	--	--	--	--	--
<i>Strep. pneumoniae</i>	8	75	--	--	100	75	75	75	--	--	100	--	88	88	--	63	--	--	100	88	--	--	100	75	88	100
<i>Enterococcus faecalis</i>	147	100	100	--	--	R	R	--	--	81	90	--	R	R	99	--	100	100	100	27	100	67	--	--	--	--
<i>Enterococcus faecium</i>	12	8	17	--	--	R	R	--	--	17	17	--	R	R	67	--	75	100	67	8	--	--	--	--	--	--
<i>Enterococcus spp.</i>	5	100	100	--	--	R	R	--	--	80	80	--	R	R	100	--	100	100	60	40	--	--	--	--	--	--

The % susceptible for each organism/antimicrobial combination was generated by including the first isolate of that organism encountered on a given patient.

R Organism has intrinsic resistance to this antimicrobial.

-- Not tested/indicated for organism.

* Antibiograms created for organisms with less than 30 total isolates have questionable statistical significance. Interpret data with caution.

(1) Oxacillin predicts susceptibility to most cephalosporins, carbapenems, and beta-lactam/beta-lactamase inhibitors.

(2) For *S. pneumoniae*: cefotaxime, ceftriaxone, and penicillin % susceptible listed is based on parenteral CSF (meningitis), parenteral non-CSF, and oral MIC breakpoints.

(3) Nitrofurantoin is reported for isolates from urine only.

(4) Linezolid is reported for isolates from non-urine sources only.

(5) For *Staphylococcus spp.*: clindamycin and erythromycin are reported for isolates from non-urine sources only.

(6) For *Staphylococcus spp.*: daptomycin is reported for isolates from non-respiratory sources only.

(7) For all *Enterococcus spp.*: ciprofloxacin, levofloxacin, and tetracycline are reported for isolates from urine only.

(8) For *E. faecalis* and *E. faecium*: gentamicin and streptomycin high-level resistance testing is reported for isolates from blood cultures only.

Physicians Laboratory Services Cumulative Antimicrobial Susceptibility Report
Hospital/Surgical Clients, All Isolates | January 2024 - December 2024

% Susceptible Gram Negative	Total*	Ampicilin	Ampicillin/Sulbactam	Piperacillin/Tazobactam	Cefazolin	Ceftazidime	Ceftazidime/Avibactam	Ceftriaxone	Cefepime	Cefoxitin	Aztreonam	Ertapenem	Imipenem	Meropenem	Amikacin	Gentamicin	Tobramycin	Ciprofloxacin(3)	Levofloxacin(3)	Trimethoprim/Sulfa	Nitrofurantoin (1)	Tetracycline(2)	Tigecycline
<i>Escherichia coli</i>	621	64	71	99	76	90	100	90	92	96	91	100	100	100	100	92	94	84	84	80	97	78	100
<i>Klebsiella pneumoniae</i>	162	R	85	99	90	93	100	93	94	94	94	100	99	100	100	97	96	95	98	86	56	82	98
<i>Proteus mirabilis</i>	91	88	91	100	79	98	100	98	99	98	99	100	--	100	100	97	97	81	90	87	R	R	--
<i>Klebsiella (Enterobacter) aerogenes</i>	35	R	R	100	R	80	100	77	94	R	91	100	74	100	100	97	97	100	100	94	23	91	100
<i>Enterobacter cloacae</i> complex	53	R	R	74	R	74	100	55	89	R	72	81	89	98	100	98	100	98	98	91	35	94	96
<i>Serratia marcescens</i>	16	R	R	81	R	81	100	81	88	R	88	94	--	100	100	100	100	88	100	94	R	13	88
<i>Pseudomonas aeruginosa</i>	131	R	R	98	R	99	100	R	98	R	94	R	90	98	98	86	100	92	94	R	--	R	--
<i>Acinetobacter baumannii</i>	1	R	100	--	R	100	--	100	100	R	R	R	100	100	100	100	100	100	100	100	--	100	--
<i>Citrobacter freundii</i> complex	36	R	R	97	R	86	100	78	94	R	83	97	94	100	100	92	94	97	97	86	87	83	100
<i>Morganella morganii</i>	15	R	0	100	R	93	100	93	100	93	100	100	0	100	100	93	87	87	93	80	R	R	0
<i>Providencia</i> spp.	2	R	0	100	R	100	100	100	100	100	100	100	100	100	100	50	50	50	50	50	R	R	0
<i>Salmonella</i> spp.	0	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
<i>Shigella</i> spp.	0	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
<i>Stenotrophomonas</i>	9	R	R	R	R	56	--	R	--	R	R	R	R	R	R	R	R	--	100	100	--	R	--
<i>Klebsiella oxytoca</i>	46	R	65	100	15	98	100	98	98	98	98	100	100	100	100	98	98	93	98	93	90	89	100
<i>Haemophilus influenzae</i>	15	Beta-lactamase positive: 27%																					

The % susceptible for each organism/antimicrobial combination was generated by including the first isolate of that organism encountered on a given patient.

R Organism has intrinsic resistance to this antimicrobial.

-- Not tested/indicated for organism.

* Antibigrams created for organisms with less than 30 total isolates have questionable statistical significance. Interpret data with caution.

(1) Nitrofurantoin is reported for isolates from urine only.

(2) For *A. baumannii*: tetracycline is reported for isolates from urine only.

(3) Ciprofloxacin/Levofloxacin: Enterobacterales & *P. aeruginosa* %Susceptible utilizes CLSI M100 28th Edition breakpoints.